

ON SOME RELATIVISTIC SINGULAR SURFACES

ANGELO LOINGER AND TIZIANA MARSICO

ABSTRACT. In the current relativistic literature there are misleading considerations about some singular surfaces. An accurate geometric analysis allows to settle the question. No physical meaning is attributable to the spatial regions surrounded by the above surfaces. A recent observational paper by A. Fabian *et al.* is discussed in sect. 4.

PACS 04.20 – General relativity.

1. – Levi-Civita has given a beautiful, *geometrically detailed* treatment of the Schwarzschild manifold created by a gravitating point-mass m [1]. First of all, following a Palatini's method [2], Levi-Civita makes a precise illustration of the geometrical concepts that suggest how to choose a convenient *system of coordinates* in a three-dimensional metric space V_3 endowed with a spherical symmetry around one of its points O . He institutes a one-to-one correspondence between V_3 and an auxiliary three-dimensional Euclidean space V'_3 (a *Bildraum*, according to Weyl's terminology), having a spherical symmetry around a point O' . Then, he proves that at any *ray* j' starting from O' there corresponds in V_3 a *geodesic line* j starting from O ; and that at any *spherical surface* Σ' of V'_3 , with centre at O' , there corresponds in V_3 a *geodesic sphere* Σ with centre at O . Let $d\sigma'$ be an elementary interval on Σ' drawn from one of its points Q' , and $d\sigma$ the homologous interval drawn from the corresponding point Q on Σ . If we refer the *Bildraum* V'_3 to polar coordinates r, ϑ, φ , we have

$$(1) \quad d\sigma'^2 = r^2(d\vartheta^2 + \sin^2 \vartheta d\varphi^2) \quad ; \quad r = O'Q' \quad ;$$

owing to the correspondence between V'_3 and V_3 , the coordinates ϑ and φ can be also considered as *curvilinear coordinates on* Σ . It is easy to prove that the correspondence between Σ and Σ' is a *conformal* one, *i.e.* that there exist a function $H(r)$ such that

$$(2) \quad d\sigma^2 = H^2(r) d\sigma'^2 \quad ;$$

if $R(r) := rH(r)$, the square of the line element $d\sigma$ of Σ can be written

$$(3) \quad d\sigma^2 = R^2(r) (d\vartheta^2 + \sin^2 \vartheta d\varphi^2) \quad .$$

In a second Euclidean *Bildraum*, eq. (3) gives the elementary interval on a sphere of radius $R(r)$, of which $K := 1/R^2$ is the Gaussian curvature,

and $S := 4\pi R^2$ the superficial area: two *invariant* concepts. Consequently, in the metric space V_3 the *coordinate* R has the following meaning: at any point P , $1/R^2$ represents the Gaussian curvature of the geodesic sphere Σ with centre O , passing through P . Further, since all geodesic lines starting from O cut orthogonally Σ , the interval dl of V_3 is given by

$$(4) \quad dl^2 = dg^2 + d\sigma^2 \quad ,$$

where dg is the elementary arc of one of the geodesics – arc which depends only on R :

$$(5) \quad dg = A(R) dR \quad ;$$

from which:

$$(6) \quad dl^2 = A^2(R) dR^2 + R^2 (d\vartheta^2 + \sin^2 \vartheta d\varphi^2) \quad .$$

This result holds under “the obvious condition that the coefficients of dl^2 are regular in the region round every point, except possibly the point O .”

If $ds^2 = U(R) dt^2 - dl^2$ is the interval of the Schwarzschild manifold, Einstein equations $R_{jk} = 0$, ($j, k = 0, 1, 2, 3$), tell us that ($c = G = 1$):

$$(7) \quad ds^2 = \left(1 - \frac{2m}{R(r)}\right) dt^2 - \frac{[dR(r)]^2}{\left(1 - \frac{2m}{R(r)}\right)} - R^2(r) (d\vartheta^2 + \sin^2 \vartheta d\varphi^2) \quad ;$$

if we put $R(r) \equiv r$, we obtain the *standard* (Hilbert-Droste-Weyl) form of ds^2 . For $R(r) \equiv [r^3 + (2m)^3]^{1/3}$, we have the *original* Schwarzschild form (1916). For $R(r) \equiv r + 2m$, the Brillouin form (1923). *Etc.* – By virtue of eq. (6), $1 - 2m/R(r)$ cannot be negative. Remark that Schwarzschild’s and Brillouin’s metrics satisfy the above obvious condition; the standard metric holds only for $r > 2m$.

2. – We have given a faithful summary of the first part of Levi-Civita’s argumentation, because it evidences the geometric meaning of the radial coordinate $R(r)$.

The *distance* D of a generic point $R(r) = \varrho$ from $R(r) = 2m$ is obviously given by

$$\begin{aligned} D &= \int_{2m}^{\varrho} \left[\frac{R(r)}{R(r) - 2m} \right]^{1/2} dR(r) = \\ (8) \quad &= \{R(r) [R(r) - 2m]\}^{1/2} + m \ln \frac{R(r) - m + \{R(r) [R(r) - 2m]\}^{1/2}}{m} \quad ; \end{aligned}$$

and *since* $R(r)/[R(r) - 2m] = A^2[R(r)]$, we see that $R(r) > 2m$.

Obviously, the “soft” singularity of interval (7) at $R(r) = 2m$ does not have a radial *distance* $d = 2m$ from $R(r) = 0$. The popular formula

$$(9) \quad (S/4\pi)^{1/2} \approx 3 \text{ km } \frac{\text{point-mass } m}{\text{solar mass}}$$

gives only a formal value of the radial *coordinate*. But the essential fact is that for $R(r) \leq 2m$ metric (7) loses any validity; consequently, *space region* $R(r) \leq 2m$ *does not admit of any reasonable interpretation*. In particular, formula (9) is problematic.

Some authors have asserted that when $R(r) < 2m$ the spatial coordinate $R(r)$ could acquire a time character and the temporal coordinate t could acquire a space character. But this change of stead would give a ds^2 which has *no* relation with our *static* problem – as it was irrefutably demonstrated many years ago by Brillouin [3].

2bis. – The second fundamental memoir of GR by Schwarzschild (1916) [4] ends with this remark: a sphere F of an incompressible fluid, having a *given* gravitating mass m , has a minimal value of its *physical radius* which is equal to $(9/8)(2m)$: a result with a real meaning, because it was obtained through the concordance at the surface of F between external and internal ds^2 , *in absence of any whatever singularity*.

3. – Of course, for Kerr’s metric [5] and for the metric of a gravitating mass-point with an electric charge [6] we can make considerations very similar to those of sect. **2**.

4. – A test-particle or a light-ray moving along a radial geodesic of Schwarzschild manifold cannot overcome the “barrier” at $R(r) = 2m$, owing to Hilbert *repulsive* effect [7]. A quite analogous result holds for Kerr’s metric [5] and for the metric of a charged mass-point [6].

Accordingly, *the involved singularities cannot swallow anything*. However, in the recent literature we find many assertions that the authors have detected various voracious eaters of matter and light, that they identify with BH’s – see *e.g.* the paper by Fabian *et al.* [8], which would concern a supermassive BH. This team of astrophysicists has observed a broad iron K and L line emission in the Narrow-Line Seyfert 1 Galaxy 1H0707-495, using the XMM-Newton satellite. The observational data tell that this Galaxy is spinning very rapidly and is swallowing the equivalent of two Earths per hour. Fabian *et al.* [8] think that at the galactic heart there is a supermassive BH, whose mass is estimated at about 3 to 5 million solar masses. The researchers would be tracking matter to within twice the “radius” of the hypothesized BH – *evidently, they utilize formula* [9]. According to Fabian, this BH is a messy eater, since “accretion is a very messy process because of the magnetic fields that are involved.”

Now, Wolfgang Kundt, a distinguished astrophysicist who does not object to the theoretical belief in the existence of BH's, has repeatedly emphasized that there are very realistic explanations of the observational data supposedly regarding various kinds of BH's, in particular the supermassive ones – see *e.g.* his concise *Astrophysics* [9].

He remarks that the active galactic nuclei (AGN), that are commonly thought to harbour a supermassive BH, “may, instead, simply be the high-density nuclear-burning centers of galactic disks.” Indeed “nuclear burning (of H to Fe) is almost as efficient a lamp as black-hole accretion, yielding $\lesssim 1\%$ of the rest-energy”, as it is suggested by nine reasons (pp.101, 102 of [9])! And further: “The QSO phenomenon may rather be a straightforward consequence of the spiralling-in of matter through *galactic disks* whose density approaches stellar values near their centers, giving rise to almost relativistic Kepler velocities on the innermost *Solar-System scales*, to strong magnetization and magnetic reconnections, and to nuclear burning in the disk's midplane. Magnetic reconnections create the pair plasma which is responsible for the jet phenomenon and for the hard, non thermal spectra, whereas nuclear power is used to re-eject matter at higher than SN velocities through the BLR. Active galactic nuclei may owe their extreme properties to those of their central disks.”

5. – *Two final remarks.* –*i*) The original Schwarzschild's form of the ds^2 , which corresponds to $R(r) \equiv [r^3 + (2m)^3]^{1/3}$, and Brillouin's form, which corresponds to $R(r) \equiv r + 2m$, are the *unique* maximally extended (and geodesically complete) metric forms of Schwarzschild manifold that do not impair, or partially suppress, the gravitational field of the point-mass. –*ii*) It is commonly believed (without a *rigorous* proof) that since p/c^2 gives a contribution to the mass density of any celestial body, even infinite pressure gradients cannot support a star of a sufficiently great mass against its self-attraction, with the formation of a BH as a final outcome. Now, this opinion is not right, as McVittie [10] and the present authors [11] have demonstrated with precise computations; a result which corroborates a famous conjecture by Eddington.

REFERENCES

- [1] T. Levi-Civita, *The Absolute Differential Calculus (Calculus of Tensors)* – (Dover Publications, Inc., Mineola, N.Y.) 2005, p. 408 *sqq.* Originally published: London: Blackie and Son, 1926. – See also H. Weyl, *Raum-Zeit-Materie*, Siebente Auflage (Springer-Verlag, Berlin, *etc.*) 1988, sect.**33**.
- [2] A. Palatini, *Nuovo Cimento*, **14** (1917) 12.
- [3] M. Brillouin, *Journ. Phys. Rad.*, **23** (1923) 43; an English translation in *arXiv:physics/0002009* (February 3rd, 2000). And references therein. – Notice that Brillouin denotes with R his radial coordinate, for which $0 \leq R < \infty$.
- [4] K. Schwarzschild, *Berl. Ber.*, (1916) 424; an English translation in *arXiv:physics/9912033* (December 16th, 1999).

- [5] A. Loinger and T. Marsico, *arXiv:0809.1221 v1* [physics.gen-ph] 7 Sept 2008. And references therein.
- [6] A. Loinger and T. Marsico, *arXiv:0907.2895 v1* [physics.gen-ph] 16 Jul 2009.
- [7] A. Loinger and T. Marsico, *arXiv:0904.1578 v1* [physics.gen-ph] 9 Apr 2009 – and references therein, in particular [1], [2].
- [8] A. Fabian *et al.*, *Nature*, **459** (2009) 540.
- [9] W. Kundt, *Astrophysics – A Primer* (Springer-Verlag, Berlin, *etc.*) 2001, pp. 101, 102, 111 – and references therein.
- [10] G.C. Mc Vittie, *Astrophys. J.*, **140** (1964) 401.
- [11] T. Marsico and A. Loinger, *arXiv:physics/0612160 v1* (December 16th, 2006). And references therein.

A.L. – DIPARTIMENTO DI FISICA, UNIVERSITÀ DI MILANO, VIA CELORIA, 16 - 20133 MILANO (ITALY)

T.M. – LICEO CLASSICO “G. BERCHET”, VIA DELLA COMMENDA, 26 - 20122 MILANO (ITALY)

E-mail address: `angelo.loinger@mi.infn.it`

E-mail address: `martiz64@libero.it`